| QCC Goals Scorecard - Proposed Final Scorecard ¹ | | | | | | | |
|---|---|-----------------|------|------|------|------|--|
| QCC Goal | Proposed Measure | | Year | | | | National Benchmark |
| QCC Goal | Proposed ineasure | 2008 | 2009 | 2010 | 2011 | 2012 | National Benchmark |
| | Poverty rate (below 100% FPL) | 11.3% | | | | | 13.2% |
| | | White: 11.6% | | | | | 12.3% |
| | By race/ethnicity | Black: 29.5% | | | | | 33.2% |
| | | Hispanic: 40.8% | | | | | 30.6% |
| Context and Resources | Adults (25+) with 4-year college degree | 21.7% | | | | | 17.5% |
| | | White: 22.7% | | | | | 21.1% |
| | By race/ethnicity | Black: 23.1% | | | | | 13.6% |
| | | Hispanic: 10.5% | | | | | 9.4% |
| | PCP's per 1,000 population | 1.8 | | | | | 1.2 |
| | Life expectancy | 79.8 | | | † | | 77.7 |
| | Infant mortality rate (deaths per 1,000 live births) | 4.8 | | | | | 6.9 |
| | Adults - overweight or obese | 58.0% | | | | | 63.0% |
| | By race/ethnicity | White: 55.2% | | | | | 59.6% |
| | | Black: 63.2% | | | | | 69.9% |
| | | Hispanic: 60.2% | | | | | |
| | Adults - current smokers | 16.0% | | | | | |
| | By race/ethnicity | White: 16.1% | | | | | |
| | | Black: 17.4% | | | | | + |
| Health Status | -,,, | Hispanic: 14.9% | | | | | |
| | Prevalence of diabetes | 7.20% | | | | | |
| | | White: 6.7% | | | | | |
| | By race/ethnicity | Black: 11.1% | | | | | |
| | -,,, | Hispanic: 8.6% | | | | | 62.1% 18.3% 18.7% 20.6% 15.6% 8.3% 7.10% 12.1% 15.4% |
| | New HIV/AIDS cases (per 100,000 people) | 8.4 | | | | | |
| | | White: 4.3 | | | | | 11.5 |
| | By race/ethnicity | Black: 35.8 | | | | | 83.8 |
| | | Hispanic: 22.1 | | | | | 29.4 |
| Health Outcomes | Age-adjusted death rate for accidents (unintentional injuries) per 100,000 population | 26.7 | | | | | 39.8 |
| | By race/ethnicity | White: 27.2 | | | | t | 39.2 |
| | | Black: 25.5 | | | | | 42.1 |
| | | Hispanic: 24.9 | | | | | 31.5 |
| | Death rate for breast cancer (per 100,000 women) | 23.0 | | | | | 24.0 |
| | By race/ethnicity | White: 23.2 | | | | | 23.3 |
| | | Black: 29.0 | | | | | 32.8 |
| | | Hispanic: 15.5 | | | | | 15.1 |
| | Death rate for colorectal cancer (per 100,000 population) | 17.6 | | | | | 17.5 |

| QCC Goal | | - Proposed Final Scorecard ¹ Year | | | | | National Benchmark |
|---------------------------------|--|--|------|------|------|----------|--------------------|
| QCC GOAI | Proposed Measure | 2008 | 2009 | 2010 | 2011 | 2012 | National Benchmark |
| | Rate of growth in per capita health care | | | | | | |
| | spending | | | | | | |
| | Rate of growth in spending by Medicaid, | | | | | | |
| | Medicare, commercial insurers, self-insured | | | | | | |
| | plans | | | | | | |
| 1. Reduce the cost of health | Consumer cost sharing as a precentage of | | | | | | |
| care. Reduce the annual rise in | total premiums | | | | | | |
| health care costs to no more | Employee contributions toward premiums | Individual = 28%; | | | | | |
| than the unadjusted growth in | Descritors of impatient beguitalizations for all | Family = 32%; (2009) | | | | | |
| GDP by 2012. | Percentage of inpatient hospitalizations for all | 8% (2006) | | | | | 10% (2006) |
| | payers that are estimated to have been preventable | 870 (2000) | | | | | 10% (2000) |
| | Inpatient days per 1,000 population | 688.5 (2007) | | | | | 657.3 (2007) |
| | Inpatient beds per 1,000 population | 2.5 (2007) | | | | | 2.7 (2007) |
| | Percent of total lives covered under risk- | , | | | | | ` ′ |
| | contracting arrangements | | | | | | |
| | Hospital Associated Infections (HAIs) | | | | | • | |
| | CLABSI rate | | | | | | 1.93 |
| | SSI | | | | | | 19.77 |
| | Percent of surgical patients receiving | | | | | | |
| | appropriate care to prevent complications | 90.3% | | | | | 85.3% |
| | appropriate care to prevent complications | | | | | | |
| | Serious Reportable Events (SRE's) ² | | | | | | |
| | Number of hospitals with SRE's | 65 | 64 | | | | |
| | Number of SRE's reported by hospitals | 338 | 383 | | | | |
| | Falls | 224 | 199 | | | | |
| | Surgical events | 62 | 76 | | | | |
| | Care management events | 26 | 78 | | | | |
| | Hospital Readmissions | , | | | , | , | |
| | Short-stay nursing home residents with a | | | | | | |
| | hospital readmission within 30 days | 19.5% | | | | | 20.8% |
| | | | | | | | |
| | Medicare 15- and 30-day readmissions | 19.4% | | | | | 17.5% |
| | Preventive Care | T T | | | 1 | | |
| | Children 19-25 months with recommended | 91.1% | | | | | 85.7% |
| 2. Ensure patient safety and | vaccines | | | | | | |
| effectiveness of care. | Children 3-6 years who saw their PCP in the | | | | | | |
| | past year | | | | | | |
| | Adults who did not visit the dentist in the past | 20.7% | | | | | 28.7% |
| | year | White: 20.8% | | | | | 26.5% |
| | By race/ethnicity | Black: 29.5% | | | | | 38.1% |
| | by race/ethnicity | Hispanic: 28.4% | | | | | 38.1% |
| | Children who did not visit the dentist in the | | | | | | |
| | past year | 16.20% | | | | | 21.6% |
| | pust yeur | White: 13.8% | | | | | 19.1% |
| | By race/ethnicity | Black: 18.0% | | | | | 21.7% |
| | 1 | Hispanic: 26.2% | | | | | 28.5% |
| | Women (40+) who have had a mammogram in | | | | | | |
| | the past two years | 87.3% | | | | | 79.5% |
| | , | White: 84.8% | | | | | 75.9% |
| | By race/ethnicity | Black: 86.6% | | | | | 79.3% |
| | | Hispanic: 88.6% | | | | | 73.5% |
| | Adults (50+) who have ever had a | | | | | | |
| | sigmoidoscopy or a colonoscopy | 71.4% | | | | <u> </u> | 62.2% |
| | High risk nursing home patients with pressure | 10.9% | | | | | 11 50/ |
| | sores | 10.9% | | | | | 11.5% |

| QCC Goals Scorecard - Proposed Final Scorecard ¹ | | | | | | | |
|--|--|---------------|------|------|------|------|--------------------|
| 000 000 | Proposed Measure | | Year | | | | National Benchmark |
| QCC Goal | | 2008 | 2009 | 2010 | 2011 | 2012 | National Benchmark |
| | Diabetes | | | | | | |
| | Avoidable hospitalization rate for patients with long-term diabetes complications (number of admissions per 100,000 population) | 126.0 | | | | | 124.9 |
| | population | White: 105 | | | | | |
| | By race/ethnicity | Black: 273 | | | | | N/A |
| | | Hispanic: 107 | | | | | |
| | Diabetes patients with HbA1c - Poor Blood Sugar Control | 17.5% | | | | | 28.4% |
| | Asthma | | | | | | |
| | Avoidable hospitalization rate - adult (number of admissions per 100,000 population) | 156 | | | | | 120.0 |
| | | White: 120 | | | | | |
| | By race/ethnicity | Black: 336 | | | | | N/A |
| | | Hispanic: 189 | | | | | |
| | Pediatric ED visit rate (per 100,000 population) | | | | | | 2330.6 |
| 3. Improve screening for and management of chronic illnesses | Adult ED visit rate (per 100,000 population) | | | | | | 1815.6 |
| in the community | Heart Disease | | | | | | |
| | Avoidable hospitalization rate for patients with CHF (number of admissions per 100,000 | 416 | | | | | 476.4 |
| | population) By race/ethnicity | White: 436 | | | | | |
| | | Black: 578 | | | | | N/A |
| | | Hispanic: 198 | | | | | 1.7 |
| | Cholesterol management for patients with cardiovascular conditions (Cholesterol (LDL-C) - Good Control) | 67.9% | | | | | 59.7% |
| | Persistence of beta blocker treatment 6 | | | | | | |
| | months after a heart attack | | | | | | |
| | Behavioral/Mental Health | | | | | | |
| | Children who received needed mental health care in the past year | 66.6% | | | | | 63.0% |
| | Antidepressant medication management (% of patients diagnosed with depression who filled a prescription for at least a 3-month supply of their medication) | 66.8% | | | | | 63.1% |

| QCC Goals Scorecard - Proposed Final Scorecard ¹ | | | | | | | | |
|---|--|--------------------|------|------|------|------|-----------------------------------|--|
| QCC Goal | Proposed Measure | Year | | | | | National Bouckersule | |
| | Proposed Measure | 2008 | 2009 | 2010 | 2011 | 2012 | National Benchmark | |
| | End of Life Care | | | | | | | |
| | State status in gaining endorsement from the National POLST Paradigm Task Force | Developing program | | | | | 7 states have an endorsed program | |
| 4. Develop and provide useful | Percent of health care settings with a palliative care program | | | | | | | |
| measurements of or approaches | | 50% | | | | | 53% | |
| to quality in areas of health care | Nursing Homes/SNFs | | | | | | | |
| for which current data are | Home Health Care | | | | | | | |
| inadequate or current | Care Transitions | | | | | | | |
| approaches are unsuccessful. | Follow-up after hospitalization for mental illness | | | | | | | |
| | MA skilled nursing facilities participating in the Advancing Excellence Program who use the "Advance Care Planning Module" | | | | | | | |
| through transparency. | Average monthly hits to MHCO Patients who access medical records through an online portal | 2586 | | | | | N/A | |

Footnotes

1 - Where available, data for measures is provided by race and/or ethnicity

⁻ The numbers of SREs reported by hospitals are likely to increase before they decrease, as hospitals become more comfortable with reporting this measure